

PROMOJEN

Instructions on medical use of the drug

Tradename: Promojen.

MNN: Turmeric, borneol.

Dosage form: Vaginal suppositories.

Composition: *One suppository contains:* Active substances:

turmeric oil82 mg;

borneol75 mg;

Excipients: ethanol, macrogol 400, macrogol 4000, polyoxyl (40) stearate, laurocapram .

Description: suppositories are yellowish-white, yellowish or brown, bullet- shaped.

Pharmacotherapeutic group: Other gynecological drugs.

Pharmacologic effect:

Pharmacodynamics :

Turmeric oil and borneol, which are part of the drug, are volatile substances, and therefore penetrate deep into the lesion. Having a wide spectrum of antibacterial and anti-inflammatory effects, turmeric oil and borneol increase the number of white blood cells in the peripheral vascular bed, improve the phagocytic activity of phagocytes and immunity. The active components of the drug improve blood circulation in the area of erosion, remove necrotic tissue, accelerate the renewal and regeneration of tissue damaged by inflammation. The drug inhibits and destroys tumor cells, but has no effect on normal tissue. Used against human papillomavirus, inhibits pathological tissue growth.

The active components of the drug inhibit the growth and reproduction of *Candida albicans* and *Torulopsis glabrata* . The drug inhibits the growth and reproduction of *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Streptococcus*, β - *Streptococcus*, *Micrococcus luteus*, *Neisseria gonorrhoeae* , *Escherichia coli*, *Pseudomonas aeruginosa*, *Bacillus cereus*, *Corynebacterium Vaginale*, *Corynebacterium diphtheroides*, *Bacteroides fragilis*, *Pepto Streptococcus*, *Bacillus typhi*, *Vibrio cholerae*.

Effective against pathogens of urogenital trichomoniasis and mycoplasmosis.

Indications for use:

- cervical erosion;
- cervicitis;
- vaginitis (including fungal);
- vaginosis (including bacterial);
- senile colpitis;
- papillomavirus infection;
- in order to prepare for treatment with physical methods: laser therapy, electrocoagulation, cryodestruction and radio wave surgery.

Contraindications:

- Girls under 14 years old;
- Pregnancy up to 12 weeks.

Directions for use and dosage:

Vaginally. Immediately before going to bed, after washing your hands, in a lying position, place the suppository on the tip of your finger and insert it deep into the vagina (posterior fornix). Use one suppository per application. The course of treatment is 14-15 days.

When treating papillomavirus infection, 1–2 courses are required, in some cases the duration of treatment is 2–3 courses. If necessary, the course of treatment should be repeated.

Suppositories can be used to prepare for treatment with physical methods such as laser therapy, electrocoagulation, cryodestruction and radio wave surgery. When using the drug, complications are reduced and tissue regeneration time is reduced.

When treating fungal colpitis, 1–2 courses of treatment are necessary. If necessary, the course of treatment should be repeated.

Treatment of bacterial vaginosis should continue for 1-2 courses, then, for the purpose of prevention, 1-2 suppositories every week. If necessary, the course of treatment should be repeated.

Side effects:

Rarely, after administration of the drug, body temperature temporarily increases, a burning sensation, pain, itching, swelling, rash and allergies appear; in older women or patients with low estrogen levels, chills appear, in which case the dosage is reduced or the drug is stopped. After stopping the use of the drug, these unwanted reactions go away on their own.

Overdose: not detected.

Special instructions and precautions:

Patients with a predisposition to allergic reactions should use the drug with caution. When using the drug, there is a feeling of cooling in the vagina.

Interaction with other drugs:

No data.

Release form:

Vaginal suppositories No 15, with instructions for medical use in a cardboard box.

Storage conditions:

Store in a dry place, protected from light, at a temperature not exceeding 25 °C.

Keep out of the reach of children!

Best before date:

2 years. Do not use after the expiration date.

Conditions of release:

By doctor's prescription.

Made for:

MAXX-PHARM LTD.

London, Great Britain

